


1. The following is a graph of the function  $f(x)$ .



- (a) (8 points) Compute the following limits if they exist. If they do not exist, explain why.

Note  
 ~~$\lim \neq \frac{d}{dx}$~~

$\lim_{x \rightarrow -8} f(x)$	$\lim_{x \rightarrow -5} f(x)$	$\lim_{x \rightarrow 0} f(x)$	$\lim_{x \rightarrow 4} f(x)$
-4	5	DNE	-∞ (if DNE)

- (b) (2 points) Find all  $x$  values where  $f'(x) = 0$ .

$x = -8$   $\leftarrow$  (+1)      (+1 for no extras)

- (c) (3 points) On which intervals is  $f'(x) > 0$ ?

$-8 < x < -5, \quad -4 < x < 6$

- (d) (3 points) At which  $x$  coordinates is  $f$  continuous but *not* differentiable?

$x = -5$   $\leftarrow$  (+1)       $x = 6$   $\leftarrow$  (+1)      (+1 no extras)

- (e) (2 points) Give the equations of any vertical asymptotes.

$x = 4$

- (f) (3 points) Compute the following limit.

$$\lim_{x \rightarrow -2} \frac{f(x) - f(-2)}{x + 2} \therefore f'(-2) = -1$$

- (g) (3 points) What is  $f''(8)$ ?

0

2. A coin is dropped from the top of the Empire State Building, from a height of 400m. Its height  $t$  seconds after being dropped is given by the equation  $h(t) = 400 - 4.9t^2$ .

(a) (3 points) Find the velocity after it has been falling for 2 seconds.

$$h'(t) = -9.8t$$

$$v(2) = -9.8 \cdot 2 = -19.6 \frac{m}{s}$$

1/2  
for  
sign  
errors

(b) (3 points) When is the coin falling at 50 m/s?

$$-9.8t = -50$$

$$t = \frac{-50}{-9.8} \approx 5.102 \text{ sec.}$$

(c) (3 points) Find the acceleration of the coin in  $m/s^2$ .

$$a = h''(t) = -9.8 \frac{m}{s^2}$$

(d) (6 points) What is the velocity of the coin when it hits the ground?

① When does  $h(t) = 0$ ?

$$4.9t^2 = 400$$

$$t^2 = 81.63$$

3 points

$$t = \underline{\underline{9.04 \text{ seconds}}}$$

②

$$v(9.04) = -9.8(9.04)$$

$$= 88.54 \frac{m}{s}$$

$$x(t) = t - \sin t$$

$$y(t) = 1 - \cos t$$

(3)

(a)

$$\text{PT } x(\frac{\pi}{3}) = \frac{\pi}{3} - \sin\left(\frac{\pi}{3}\right) = \frac{\pi}{3} - \frac{\sqrt{3}}{2}$$

$$y\left(\frac{\pi}{3}\right) = 1 - \cos\left(\frac{\pi}{3}\right) = 1 - \frac{1}{2} = \frac{1}{2}$$

slope

$$x'(t) = 1 - \cos t \quad x'\left(\frac{\pi}{3}\right) = 1 - \frac{1}{2} = \frac{1}{2}$$

$$y'(t) = \sin t \quad y'\left(\frac{\pi}{3}\right) = \frac{\sqrt{3}}{2}$$

$$m = \frac{y'}{x'} = \frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}} = \sqrt{3}$$

Line  $y = \sqrt{3}\left(x - \frac{\pi}{3} + \frac{\sqrt{3}}{2}\right) + \frac{1}{2}$

(b)

No L? Max y

Note  $x'(t) = 0$

$$\cos t = 1 \Rightarrow t = 0 + 2\pi \cdot k$$

$$y'(t) = 0$$

$$\Leftrightarrow \sin t = 0 \Rightarrow t = 0 + \pi k$$

Therefore Horizontal (a)  $t = \pi + 2\pi k$

Vertical?

$$\lim_{t \rightarrow 0} \frac{\sin t}{1 - \cos t} \cdot \frac{1 + \cos t}{1 + \cos t} = \frac{\sin t + \sin t \cos t}{\sin^2 t} = \frac{\sin t}{\sin^2 t} \cdot \frac{1 + \cos t}{\sin t} = \frac{1 + \cos t}{\sin t}$$

$t = 0 + 2\pi k$

DO!

Plugging in #s? Max = 1

4. Compute the following limits. Indicate to me which one you skip. Show and justify all steps! If the limit does not exist explain why.

(a) (5 points)

$$\lim_{x \rightarrow \infty} \frac{\sqrt{16x^2 - 24x} - 4x}{\sqrt{16x^2 - 24x} + 4x}$$

$$= \lim_{x \rightarrow \infty} \frac{16x^2 - 24x + 16x^2}{\sqrt{16x^2 - 24x} + 4x} = \lim_{x \rightarrow \infty} \frac{-24x}{\sqrt{16x^2 - 24x} + 4x}$$

$$= \lim_{x \rightarrow \infty} \frac{-24}{\sqrt{16 - \frac{24}{x}} + 4} = \frac{-24}{4 + 4} = \underline{-3}$$

(b) (5 points) (Hint: Compute the limit from each side)

$$\lim_{x \rightarrow -1^-} \frac{|x+1|}{x+1}$$

$$\lim_{x \rightarrow -1^-} \frac{-(x+1)}{x+1}$$

$$\lim_{x \rightarrow -1^-} -1 = -1$$

$$\lim_{x \rightarrow -1^+} \frac{|x+1|}{x+1}$$

$$\lim_{x \rightarrow -1^+} \frac{x+1}{x+1}$$

$$\lim_{x \rightarrow -1^+} +1 = +1$$

| 1 ≠ -1  
So DNE

(c) (5 points)

$$\lim_{z \rightarrow 0} \frac{(2+z)^8 - 2^8}{z} = f'(2) \text{ where } f(x) = x^8$$

$$\text{Thus } f'(x) = 8x^7$$

$$f'(2) = 8 \cdot 2^7 = 2^{10} = 1024$$

\* Bonus ~~+1~~ to those who noticed this

L'Hopital  
Max = 1

5. Compute the  $\frac{dy}{dx}$  for 5 of the following curves. Indicate to me which one you skip. Show and justify all steps! Once computed, you do not need to simplify the derivative.

(a) (5 points)

$$f(x) = \sqrt[10]{x^5 + x^4 + x^3 + x^2 + x + 1} = (x^5 + x^4 + x^3 + x^2 + x + 1)^{\frac{1}{10}}$$

$$f'(x) = \frac{1}{10}(x^5 + x^4 + x^3 + x^2 + x + 1)^{\frac{-9}{10}} \cdot (5x^4 + 4x^3 + 3x^2 + 2x + 1)$$

(b) (5 points)

$$x^3 - e^{xy} + \sin(y) = 13.$$

Implicit:  $3x^2 - e^{xy}y - e^{xy}xy' + \cos y \cdot y' = 0$

$$y'(\cos y - e^{xy}) = e^{xy}y - 3x^2$$

$$y' = \frac{e^{xy}y - 3x^2}{\cos y - e^{xy}x}$$

(c) (5 points)

Product

$$y = \arctan(x) \ln(1/x^2).$$

$$y' = \frac{1}{1+x^2} \ln(1/x) + \arctan x \left( \frac{1}{1/x^2} \cdot \frac{-2}{x^3} \right)$$

+ 2  
7

+ 3  
7

in 55  
chain?

- 5


(d) (5 points)

CHAIN

$$h(x) = 3^{\ln(\sin(x^2))}$$

$$h'(x) = \ln 3 \cdot 3^{\ln(\sin(x^2))} \cdot \left( \frac{1}{\sin(x^2)} \right) \cdot (\cos x^2) \cdot (2x)$$

No Ch.n?



(e) (5 points)

Log Diff

$$l(t) = \frac{3x^2\sqrt{2x-5}}{\sin(x)\cos(x)}$$

$$\ln(y) = \ln(3x^2\sqrt{2x-5}) - \ln(\sin x \cos x)$$

$$= \ln 3 + 2\ln(x) + \frac{1}{2}\ln(2x-5)$$

$$-\ln \sin x - \ln \cos x$$

$$\frac{g'}{y} = \frac{2}{x} + \frac{1}{2x-5} - \frac{\cos x}{\sin x} + \frac{\sin x}{\cos x} \quad \left| \begin{array}{l} g' = y' \\ y = g(t) \end{array} \right.$$

(f) (5 points)

$$y = \sin(x)^{\cos(x)}$$

log diff

$$\ln y = \cos x \ln \sin x$$

$$\frac{g'}{y} = -\sin x \ln \sin x + \cos x \cdot \frac{\cos}{\sin x}$$

$$y' = \sin x^{\cos x} \left( \frac{\cos^2 x}{\sin x} - \sin x \ln \sin x \right).$$

No log? No credit  
X X X X X